




BRCA1 MEN



The medical management guidelines from the National Comprehensive Cancer Network (NCCN) for patients with a BRCA1 pathogenic / likely pathogenic variant are listed in this document.

This overview is for informational purposes and does not constitute a personalised recommendation. **Recommended options may vary based on your personal and family history. Access to some options may also vary from one medical center to another.** The specific references should be consulted for more details before developing a treatment plan.

In addition, the information available on hereditary cancer susceptibility genes is constantly evolving and **it is recommended to check this information annually as the management guidelines may change in the future.**



References:

Daly M et coll. NCCN Clinical Practice Guidelines in Oncology: Genetic/Familial High-Risk Assessment: Breast, Ovarian and Pancreatic. Version 2.2021. November 20, 2020. <http://www.nccn.org>

Petrucelli N, Daly MB, Pal T. BRCA1- and BRCA2-Associated Hereditary Breast and Ovarian Cancer. 1998 Sep 4 [Updated 2016 Dec 15]. In : Pagon RA, Adam MP, Ardinger HH, et al., editors. GeneReviews® [Internet]. Seattle (WA) : University of Washington, Seattle; 1993-2017. <https://www.ncbi.nlm.nih.gov/books/NBK1247>

Canto MI, Harinck F, Hruban RH, et coll. International Cancer of the Pancreas Screening (CAPS) Consortium summit on the management of patients with increased risk for familial pancreatic cancer. Gut 2013 ; 62:339-347.



BRCA1 MEN



BREAST CANCER

PROSTATE CANCER

SCREENING

STARTING AT AGE 35

- MONTHLY BREAST SELF-EXAMINATION.
 - ◊ PROMPTLY REPORT CHANGES TO A HEALTHCARE PROVIDER.
- CLINICAL BREAST EXAMINATION BY A PHYSICIAN, EVERY 12 MONTHS

IN MEN WITH GYNECOMASTIA (INCREASED AMOUNT OF BREAST TISSUE)

STARTING AT AGE 50

- OR 10 YEARS EARLIER THAN THE YOUNGEST MALE BREAST CANCER DIAGNOSIS IN THE FAMILY
- CONSIDER MAMMOGRAM EVERY 12 MONTHS.

35

50

40

STARTING AT AGE 40

- CONSIDER PROSTATE CANCER SCREENING (RECTAL EXAM OF THE PROSTATE AND PSA BLOOD TEST) EVERY 12 MONTHS

PANCREATIC CANCER

SCREENING

CURRENTLY, SCREENING FOR PANCREATIC CANCER IS NOT RECOMMENDED IN ABSENCE OF A FAMILY HISTORY OF PANCREATIC CANCER.

FOR PATHOGENIC/LIKELY PATHOGENIC VARIANT CARRIERS **WITH ≥1** FIRST-DEGREE RELATIVE (PARENT, CHILD OR SIBLING) OR SECOND-DEGREE RELATIVE (GRAND-PARENT, AUNT OR UNCLE, NIECE OR NEPHEW) (ON THE SAME SIDE OF THE FAMILY) DIAGNOSED WITH PANCREATIC CANCER:

STARTING AT AGE 50

OR 10 YEARS BEFORE THE EARLIEST PANCREATIC CANCER IN THE FAMILY

- CONSIDER SCREENING WITH MRI/MAGNETIC RESONANCE CHOLANGIOPANCREATOGRAPHY (MRCP) AND/OR ENDOSCOPIC ULTRASONOGRAPHY (EUS), IN AN EXPERIENCED CENTER, IDEALLY UNDER RESEARCH PROTOCOL.